

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

REGION 6 1445 ROSS AVENUE, SUITE 1200 DALLAS, TX 75202-2733

October 16, 2015



RE: Wilcox Oil Company Superfund Site - Residential Data Results

Dear Mr. and Mrs.(b)

This letter provides you with a summary of the soil data results for your property. Based on our review of the data, soil samples collected from your property did not exceed levels considered acceptable by EPA for short-term exposures. Although some concentrations exceeded their Residential Screening Level (RSL) for soil, this is merely an indication that further evaluation may be needed. The concentrations that exceeded the RSL are expected to fall within the accepted range established by EPA for long term exposures.

Table 1 presents the results for those chemicals detected in the soil and exceeding the RSL. All other data concentrations fell below the RSL. One metal and several organic chemicals were detected as levels that exceed the Residential Soil Screening Levels (RSL). Arsenic was detected in every sample with the highest result being 6.5 milligrams per kilogram (mg/kg) at a depth of 2 - 6 inches below ground surface (bgs). Published reports have documented the background range of arsenic concentrations for Oklahoma as being 0.6 to 21 mg/kg. The highest result of 6.5 mg/kg falls within the range of background concentrations. Several Polycyclic Aromatic Hydrocarbons and Benzene were detected as presented below.

	Table 1											
Grid	Depth	Chemical	Concentration mg/kg	Screening Level mg/kg								
Grid 1	0 – 2 inches bgs	Benzo(a)pyrene	0.02	0.016								
Grid 2	0 – 2 inches bgs	Benzo(a)pyrene	0.018	0.016								
	2 – 6 inches bgs	Benzo(a)pyrene	0.02	0.016								
	12 – 24 inches bgs	Benzo(a)anthracene Benzo(a)pyrene Benzo(b)fluoranthene Indeno(1,2,3-cd)pyrene	1.6 0.94 0.51 0.47	0.16 0.016 0.16 0.16								
	12 – 24 inches bgs Duplicate	Benzo(a)pyrene	0.073	0.016								
Grid 4	6 – 12 inches bgs	Benzo(a)pyrene	0.017	0.016								

The Residential Soil Screening Levels represent conservative levels developed using risk assessment guidance from the EPA Superfund program. RSLs are considered by the Agency to be protective for humans (including sensitive groups) over a lifetime of exposure. RSLs are not cleanup standards and are used for site "screening" to help identify areas, contaminants, and



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

REGION 6 1445 ROSS AVENUE, SUITE 1200 DALLAS, TX 75202-2733

October 16, 2015



RE: Wilcox Oil Company Superfund Site – Residential Data Results

Dear Mr. and Mrs.(b)

This letter provides you with a summary of the soil data results for your property. Based on our review of the data, soil samples collected from your property did not exceed levels considered acceptable by EPA for short-term exposures. Although some concentrations exceeded their Residential Screening Level (RSL) for soil, this is merely an indication that further evaluation may be needed. The concentrations that exceeded the RSL are expected to fall within the accepted range established by EPA for long term exposures.

Table 1 presents the results for those chemicals detected in the soil and exceeding the RSL. All other data concentrations fell below the RSL. One metal and several organic chemicals were detected as levels that exceed the Residential Soil Screening Levels (RSL). Arsenic was detected in every sample with the highest result being 6.5 milligrams per kilogram (mg/kg) at a depth of 2 - 6 inches below ground surface (bgs). Published reports have documented the background range of arsenic concentrations for Oklahoma as being 0.6 to 21 mg/kg. The highest result of 6.5 mg/kg falls within the range of background concentrations. Several Polycyclic Aromatic Hydrocarbons and Benzene were detected as presented below.

Table 1										
Grid	Depth	Chemical	Concentration mg/kg	Screening Level mg/kg						
Grid 1	0-2 inches bgs	Benzo(a)pyrene	0.02	0.016						
Grid 2	0-2 inches bgs	Benzo(a)pyrene	0.018	0.016						
	2 – 6 inches bgs	Benzo(a)pyrene	0.02	0.016						
	12 – 24 inches bgs	Benzo(a)anthracene	1.6	0.16						
		Benzo(a)pyrene	0.94	0.016						
		Benzo(b)fluoranthene	0.51	0.16						
		Indeno(1,2,3-cd)pyrene	0.47	0.16						
	12 – 24 inches bgs Duplicate	Benzo(a)pyrene	0.073	0.016						
Grid 4	6 – 12 inches bgs	Benzo(a)pyrene	0.017	0.016						

The Residential Soil Screening Levels represent conservative levels developed using risk assessment guidance from the EPA Superfund program. RSLs are considered by the Agency to be protective for humans (including sensitive groups) over a lifetime of exposure. RSLs are not cleanup standards and are used for site "screening" to help identify areas, contaminants, and

conditions that may require further investigation at a particular site. Generally, at sites where contaminant concentrations fall below RSLs, no further action or study is warranted under the Superfund program, so long as the exposure assumptions at a site match those used to develop the RSL. Chemical concentrations above the RSL do not automatically trigger a response action; however, exceeding the RSL suggests that further evaluation of the potential risks by site contaminants is appropriate. In addition, further evaluation of the potential risks related to site contaminants that were detected in the soil but have no RSL is appropriate. As we move forward with our site investigation, we will be evaluating these contaminants in more detail through a Site-specific Human Health Risk Assessment to determine the potential risks these contaminants may pose. For additional information related to Arsenic and Polycyclic Aromatic Hydrocarbons, please see the enclosed contaminant profiles developed by the Agency for Toxic Substances and Disease Registry (ATSDR).

If you are concerned, please consider some of these options while we evaluate these contaminants in more detail.

- Ensure proper hygiene, especially frequent hand washing.
- Soil should be thoroughly shaken off clothes and footwear, before entering homes.
- Use a dust mask when mowing.
- Limit outside digging and soil moving activities.

We would like to follow-up with you during the week of October 26, 2015, to discuss the data presented in this letter and answer any questions that you may have. We will contact you by phone to schedule a time that fits your schedule. In the interim, should you want to talk with us, please contact me at 214-665-8143, or 1-800-533-3508, or contact Todd Downham, Oklahoma Department of Environmental Quality at 405-702-5136.

Sincerely,

Katrina Higgins-Coltrain, Remedial Project Manager

LA/OK/NM Section

Enclosure 1: Understanding the Soil Collection Process


The following summary is provided to help you better understand how we sampled your property. The figure provided as Enclosure 2 shows that your property was divided into several sample grids. As shown in Figure 2, each Grid has five sample locations. From each of these five locations, a 24 inch long soil core was taken. The soil core was divided into depth intervals of 0-2 inches below ground surface (bgs), 2–6 inches bgs, 6-12 inches bgs, and 12-24 inches bgs. The soil from each depth interval was combined together to produce one sample representing that depth interval. For example, soil from 0–2 inches bgs was collected from each of the five sample locations and then combined into one sample that represents the 0-2 inch depth interval for that Grid. This process was used for each depth interval and results in a total of 4 soil samples for each Grid. One sample represents the 0–2 inches bgs, the second represents 2–6 inches bgs, the third represents 6–12 inches bgs, and the fourth represents 12–24 inches bgs. Figure 2 shows an example of the sample process. If there is no sample result for a depth interval, then we were not able to collect that sample due to little soil recovery or the core could not be pushed deeper because of rock. The table provided in Enclosure 3 summarizes the data results for all chemicals detected in the soil and provides the full list of chemicals that were analyzed. If a chemical is not summarized, then that chemical was not detected in the soil.

O-2 in sample

Figure 2: Example of Soil Collection Process

Sample location

Depth	Depths for each Sample Location								
	0 – 2 inches below ground surface (bgs)								
	2 – 6 inches bgs								
	6 – 12 inches bgs								
	12 – 24 inches bgs								



			EPA	Property 011				
			Residential	- •	Cuid 1	Cuid 1	C 1	Cuid 1
				Sample ID	Grid 1	Grid 1	Grid 1	Grid 1
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Metals								
Aluminum	7429-90-5	mg/kg	77000		7560	5470 J	7500 J	6930 J
Arsenic	7440-38-2	mg/kg	0.68		2.6 J	1.2	2.9	2.1
Barium	7440-39-3	mg/kg	15000		73.4	48.1 J	109 J	78.6 J
Beryllium	7440-41-7	mg/kg	160		0.54	0.46 U	0.76	0.7
Calcium	7440-70-2	mg/kg	NP		3160 J	2860	6470	1520
Chromium	7440-47-3	mg/kg	NP		14.8 J	9	24	18.2
Cobalt	7440-48-4	mg/kg	23		4.6	2	5.2	5.2
Copper	7440-50-8	mg/kg	3100		8.6 J	4.9	9.9	6.6
Iron	7439-89-6	mg/kg	55000		14300	9500 J	12200 J	10300 J
Lead	7439-92-1	mg/kg	400		24	6.4	9.5	7.9
Magnesium	7439-95-4	mg/kg	NP		1020 J	835	951	939
Manganese	7439-96-5	mg/kg	1800		241 J	227	134	79.7
Mercury	7439-97-6	mg/kg	9.4		0.01 LJ	0.0023 LJ	0.0037 LJ	0.11 U
Nickel	7440-02-0	mg/kg	1500		8.8	4.9	14.7	10.9
Potassium	7440-09-7	mg/kg	NP		1130	760	739	642
Selenium	7782-49-2	mg/kg	390		2.3 UJ	0.3 LJ	2.8 U	2 U
Vanadium	7440-62-2	mg/kg	390		23.1	14.6	36.6	23.7
Zinc	7440-66-6	mg/kg	23000		61.6 J	19 J	21.9 J	15.4 J



			EPA	Property 011				
			Residential	Sample ID	Grid 1	Grid 1	Grid 1	Grid 1
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Polychlorinated Biphenyls (No	ne Detected)							
Pesticides								
4,4'-DDD	72-54-8	mg/kg	2.3		0.0042 U	0.002 LJ	0.00047 LJ	0.00024 LJ
4,4'-DDE	72-55-9	mg/kg	2		0.00063 LJ	0.00051 LJ	0.00051 LJ	0.00019 LJ
4,4'-DDT	50-29-3	mg/kg	1.9		0.00052 LJ	0.00085 LJ	0.00041 LJ	0.00033 LJ
Aldrin	309-00-2	mg/kg	0.039		0.0022 U	0.00017 LJ	0.00029 LJ	0.0021 U
alpha-BHC	319-84-6	mg/kg	0.086		0.0022 U	0.0022 U	0.002 U	0.0021 U
alpha-Chlordane	5103-71-9	mg/kg	NP		0.0022 U	0.00055 LJ	0.002 U	0.0021 U
beta-BHC	319-85-7	mg/kg	0.3		0.00002 LJ	0.00049 LJ	0.00023 LJ	0.00013 LJ
delta-BHC	319-86-8	mg/kg	NP		0.00022 LJ	0.0022 U	0.002 U	0.0021 U
Dieldrin	60-57-1	mg/kg	0.034		0.0015 LJ	0.0043 U	0.0039 U	0.0041 U
Endosulfan I	959-98-8	mg/kg	NP		0.0022 U	0.0022 U	0.002 U	0.0021 U
Endosulfan II	33213-65-9	mg/kg	NP		0.00076 LJ	0.0043 U	0.00016 LJ	0.000065 LJ
Endosulfan sulfate	1031-07-8	mg/kg	NP		0.0042 U	0.0043 U	0.0039 U	0.0041 U
Endrin	72-20-8	mg/kg	19		0.0042 U	0.000058 LJ	0.0039 U	0.0002 LJ
Endrin aldehyde	7421-93-4	mg/kg	NP		0.0042 U	0.0043 U	0.0039 U	0.0041 U
Endrin ketone	53494-70-5	mg/kg	NP		0.00093 LJ	0.0012 LJ	0.0013 LJ	0.00063 LJ
gamma-BHC (Lindane)	58-89-9	mg/kg	0.57		0.00027 LJ	0.0022 U	0.002 U	0.0021 U
gamma-Chlordane	5103-74-2	mg/kg	NP		0.00068 LJ	0.0012 LJ	0.00068 LJ	0.000069 LJ
Heptachlor	76-44-8	mg/kg	0.13		0.0003 LJ	0.0022 U	0.002 U	0.00015 LJ
Heptachlor epoxide	1024-57-3	mg/kg	0.07		0.00065 LJ	0.0022 U	0.002 U	0.0021 U
Methoxychlor	72-43-5	mg/kg	320		0.022 U	0.022 U	0.02 U	0.021 U



			EPA	Property 011				
			Residential	Sample ID	Grid 1	Grid 1	Grid 1	Grid 1
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	_	6/2/2015	6/2/2015	6/2/2015
Semi-volatile Organic Compou	ınds							
Bis(2-ethylhexyl)phthalate	117-81-7	mg/kg	39		0.22 U	0.22 U	0.2 U	0.21 U
Pyrene	129-00-0	mg/kg	1800		NA	NA	NA	NA
Semi-volatile Organic Compou	ınds (SIM)							
2-Methylnaphthalene	91-57-6	mg/kg	240		0.0036 LJ	0.0053 LJ	0.0078 U	0.004 U
Acenaphthene	83-32-9	mg/kg	3600		0.0084 U	0.0085 U	0.0078 U	0.004 U
Acenaphthylene	208-96-8	mg/kg	NP		0.0034 LJ	0.003 LJ	0.0021 LJ	0.004 U
Anthracene	120-12-7	mg/kg	18000		0.0074 LJ	0.0098	0.0054 LJ	0.0016 LJ
Benzo(a)anthracene	56-55-3	mg/kg	0.16		0.026	0.02	0.017	0.0073
Benzo(a)pyrene	50-32-8	mg/kg	0.016		0.02	0.015	0.014	0.0052
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16		0.031	0.024	0.021	0.0095
Benzo(g,h,i)perylene	191-24-2	mg/kg	NP		0.0057 LJ	0.016	0.016	0.0028 LJ
Benzo(k)fluoranthene	207-08-9	mg/kg	1.6		0.0074 LJ	0.0058 LJ	0.0052 LJ	0.0015 LJ
Chrysene	218-01-9	mg/kg	16		0.047	0.036	0.037	0.025
Dibenzo(a,h)anthracene	53-70-3	mg/kg	0.016		0.0049 LJ	0.0048 LJ	0.0049 LJ	0.0023 LJ
Fluoranthene	206-44-0	mg/kg	2400		0.032	0.022	0.015	0.0053
Fluorene	86-73-7	mg/kg	2400		0.0084 U	0.0027 LJ	0.0078 U	0.004 U
Indeno(1,2,3-cd)pyrene	193-39-5	mg/kg	0.16		0.014	0.015	0.014	0.005
Naphthalene	91-20-3	mg/kg	3.8		0.0022 LJ	0.0019 LJ	0.0078 U	0.004 U
Phenanthrene	85-01-8	mg/kg	NP		0.023	0.022	0.012	0.0068
Pyrene	129-00-0	mg/kg	1800		0.039	0.061	0.024	0.012



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	-	Grid 1 0-2in bgs 6/1/2015	Grid 1 2-6in bgs 6/2/2015	Grid 1 6-12in bgs 6/2/2015	Grid 1 12-24in bgs 6/2/2015
Volatile Organic Compounds								
2-Butanone	78-93-3	mg/kg	27000		0.013 U	0.011 U	0.0074 LJ	0.0095 U
Acetone	67-64-1	mg/kg	61000		0.0021 LJ	0.0074 LJ	0.0098 LJ	0.007 LJ
Ethylbenzene	100-41-4	mg/kg	5.8		0.0063 U	0.0053 U	0.0054 U	0.0047 U
m,p-Xylene	108-38-3/106-42-3	mg/kg	550		0.0063 U	0.00027 LJ	0.0054 U	0.0047 U
o-Xylene	95-47-6	mg/kg	650		0.0063 U	0.0053 U	0.0054 U	0.0047 U
Toluene	108-88-3	mg/kg	4900		0.0063 U	0.0013 LJ	0.0026 LJ	0.0014 LJ

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL NA - Not Analyzed

Yellow highlighted cells exceed RSL NP - Not Published
U - Not Detected at reported quantitation limit J - Estimated Value

L - Reported concentration is below the CRQL R - Unusable

mg/kg-milligrams per kilogram SIM-selected ion monitoring

BOLD-indicates a detected concentration in-inches

bgs-below ground surface



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	Property 011 Sample ID Depth Date	Grid 2 0-2in bgs	Grid 2 2-6in bgs 6/2/2015	Grid 2 6-12in bgs 6/2/2015	Grid 2 12-24in bgs 6/2/2015	Grid 2- Duplicate 12-24in bgs 6/2/2015
Metals									
Aluminum	7429-90-5	mg/kg	77000		3690	3310 J	3090 J	3330 J	4870 J
Arsenic	7440-38-2	mg/kg	0.68		2 J	6.5	1	0.99	1.7
Barium	7440-39-3	mg/kg	15000		49.8	45.6 J	45 J	29.7 J	41.1 J
Beryllium	7440-41-7	mg/kg	160		0.6 U	0.57	0.47 U	0.39 U	0.44 U
Calcium	7440-70-2	mg/kg	NP		3170 J	2050	1540	515 U	587
Chromium	7440-47-3	mg/kg	NP		10.2 J	25.9	10	8.3	10.6
Cobalt	7440-48-4	mg/kg	23		3	4.5	2.7	1.6	2.2
Copper	7440-50-8	mg/kg	3100		5.8 J	8.3	5.9	4.7	5.5
Iron	7439-89-6	mg/kg	55000		6240	6820 J	6780 J	5210 J	8490 J
Lead	7439-92-1	mg/kg	400		23.7	17.3	5.7	2.6	4.6
Magnesium	7439-95-4	mg/kg	NP		1170 J	619	392 LJ	300 LJ	486 LJ
Manganese	7439-96-5	mg/kg	1800		244 J	157	161	23.1	57.5
Mercury	7439-97-6	mg/kg	9.4		0.012 LJ	0.0031 LJ	0.11 U	0.098 U	0.1 U
Nickel	7440-02-0	mg/kg	1500		5.9	7.7	5.8	4.2	6.6
Potassium	7440-09-7	mg/kg	NP		691	571	464	515 U	618
Selenium	7782-49-2	mg/kg	390		3 UJ	2.4 U	2.3 U	1.9 U	2.2 U
Vanadium	7440-62-2	mg/kg	390		14.3	20.1	15.3	12	17.7
Zinc	7440-66-6	mg/kg	23000		22.8 J	20.5 J	11.5 J	10.7 J	13.9 J



			EPA	Property 011					Grid 2-
			Residential	Sample ID	Grid 2	Grid 2	Grid 2	Grid 2	Duplicate
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015	6/2/2015
Polychlorinated Biphenyls (No	ne Detected)								
Pesticides									
4,4'-DDD	72-54-8	mg/kg	2.3		0.0045 LJ	0.001 LJ	0.00043 LJ	0.0017 LJ	0.0019 LJ
4,4'-DDE	72-55-9	mg/kg	2		0.00025 LJ	0.00059 LJ	0.0037 U	0.01 LJ	0.00039 LJ
4,4'-DDT	50-29-3	mg/kg	1.9		0.0016 LJ	0.00072 LJ	0.0002 LJ	0.0086 LJ	0.00074 LJ
Aldrin	309-00-2	mg/kg	0.039		0.01 U	0.00036 LJ	0.00029 LJ	0.0098 U	0.00011 LJ
alpha-BHC	319-84-6	mg/kg	0.086		0.00043 LJ	0.0021 U	0.0019 U	0.0098 U	0.002 U
alpha-Chlordane	5103-71-9	mg/kg	NP		0.01 U	0.00032 LJ	0.0019 U	0.0098 U	0.002 U
beta-BHC	319-85-7	mg/kg	0.3		0.00014 LJ	0.00026 LJ	0.00026 LJ	0.0098 U	0.00031 LJ
delta-BHC	319-86-8	mg/kg	NP		0.01 U	0.0021 U	0.0019 U	0.0098 U	0.002 U
Dieldrin	60-57-1	mg/kg	0.034		0.0011 LJ	0.004 U	0.0037 U	0.019 U	0.0038 U
Endosulfan I	959-98-8	mg/kg	NP		0.01 U	0.0021 U	0.0019 U	0.0034 LJ	0.00031 LJ
Endosulfan II	33213-65-9	mg/kg	NP		0.002 LJ	0.000092 LJ	0.000064 LJ	0.019 U	0.0038 U
Endosulfan sulfate	1031-07-8	mg/kg	NP		0.00047 LJ	0.00023 LJ	0.000049 LJ	0.008 LJ	0.00084 LJ
Endrin	72-20-8	mg/kg	19		0.0033 LJ	0.00067 LJ	0.00054 LJ	0.0044 LJ	0.00043 LJ
Endrin aldehyde	7421-93-4	mg/kg	NP		0.02 U	0.004 U	0.0037 U	0.019 U	0.0038 U
Endrin ketone	53494-70-5	mg/kg	NP		0.02 U	0.00052 LJ	0.00014 LJ	0.016 LJ	0.0013 LJ
gamma-BHC (Lindane)	58-89-9	mg/kg	0.57		0.01 U	0.0021 U	0.0019 U	0.0098 U	0.002 U
gamma-Chlordane	5103-74-2	mg/kg	NP		0.0012 LJ	0.0011 LJ	0.00022 LJ	0.0028 LJ	0.00029 LJ
Heptachlor	76-44-8	mg/kg	0.13		0.01 U	0.0021 U	0.0019 U	0.0098 U	0.002 U
Heptachlor epoxide	1024-57-3	mg/kg	0.07		0.00034 LJ	0.0021 U	0.0019 U	0.0098 U	0.002 U
Methoxychlor	72-43-5	mg/kg	320		0.1 U	0.021 U	0.019 U	0.098 U	0.02 U



			EPA	Property 011					Grid 2-
			Residential	Sample ID	Grid 2	Grid 2	Grid 2	Grid 2	Duplicate
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015	6/2/2015
Semi-volatile Organic Compou	ınds								
Bis(2-ethylhexyl)phthalate	117-81-7	mg/kg	39		2.1 U	0.03 LJ	0.19 U	6 U	0.99 U
Pyrene	129-00-0	mg/kg	1800		NA	NA	NA	19	1.3
Semi-volatile Organic Compou	ınds (SIM)								
2-Methylnaphthalene	91-57-6	mg/kg	240		0.04 U	0.0017 LJ	0.0075 U	0.16 LJ	0.093 LJ
Acenaphthene	83-32-9	mg/kg	3600		0.04 U	0.0081 U	0.0075 U	0.73 LJ	0.056 LJ
Acenaphthylene	208-96-8	mg/kg	NP		0.04 U	0.0025 LJ	0.0075 U	0.77 U	0.23 U
Anthracene	120-12-7	mg/kg	18000		0.04 U	0.0063 LJ	0.0033 LJ	2.1	0.13 LJ
Benzo(a)anthracene	56-55-3	mg/kg	0.16		0.02 LJ	0.02	0.013	1.6	0.12 LJ
Benzo(a)pyrene	50-32-8	mg/kg	0.016		0.018 LJ	0.02	0.0089	0.94	0.073 LJ
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16		0.026 LJ	0.026	0.016	0.51 LJ	0.23 U
Benzo(g,h,i)perylene	191-24-2	mg/kg	NP		0.014 LJ	0.046	0.01	0.72 LJ	0.053 LJ
Benzo(k)fluoranthene	207-08-9	mg/kg	1.6		0.04 U	0.0062 LJ	0.0049 LJ	0.77 U	0.23 U
Chrysene	218-01-9	mg/kg	16		0.027 LJ	0.029	0.015	1.9	0.15 LJ
Dibenzo(a,h)anthracene	53-70-3	mg/kg	0.016		0.04 U	0.0069 LJ	0.0028 LJ	0.77 U	0.23 U
Fluoranthene	206-44-0	mg/kg	2400		0.023 LJ	0.021	0.016	1.4	0.11 LJ
Fluorene	86-73-7	mg/kg	2400		0.04 U	0.0081 U	0.0075 U	1.5	0.097 LJ
Indeno(1,2,3-cd)pyrene	193-39-5	mg/kg	0.16		0.021 LJ	0.027	0.011	0.47 LJ	0.23 U
Naphthalene	91-20-3	mg/kg	3.8		0.04 U	0.0081 U	0.0075 U	0.77 U	0.23 U
Phenanthrene	85-01-8	mg/kg	NP		0.014 LJ	0.0096	0.0032 LJ	4.8	0.35
Pyrene	129-00-0	mg/kg	1800		0.029 LJ	0.032	0.018	NA	NA



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	Property 011 Sample ID Depth Date	Grid 2	Grid 2 2-6in bgs 6/2/2015	Grid 2 6-12in bgs 6/2/2015	Grid 2 12-24in bgs 6/2/2015	Grid 2- Duplicate 12-24in bgs 6/2/2015
Volatile Organic Compounds									
2-Butanone	78-93-3	mg/kg	27000		0.015 U	0.017	0.0094 LJ	0.014 LJ	0.016 U
Acetone	67-64-1	mg/kg	61000		0.0043 LJ	0.041	0.0092 LJ	0.018	0.014 LJ
Ethylbenzene	100-41-4	mg/kg	5.8		0.0077 U	0.0061 U	0.0054 U	0.007 U	0.0081 U
m,p-Xylene	108-38-3/106-42-3	mg/kg	550		0.0077 U	0.0061 U	0.0054 U	0.007 U	0.0081 U
o-Xylene	95-47-6	mg/kg	650		0.00025 LJ	0.0061 U	0.0054 U	0.007 U	0.0081 U
Toluene	108-88-3	mg/kg	4900		0.011 J^	0.0061	0.0015 LJ	0.0024 LJ	0.00079 LJ

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL

NA - Not Analyzed NP - Not Published

Yellow highlighted cells exceed RSL

J - Estimated Value

U - Not Detected at reported quantitation limit L - Reported concentration is below the CRQL

R - Unusable

mg/kg-milligrams per kilogram

SIM-selected ion monitoring

BOLD-indicates a detected concentration

in-inches

bgs-below ground surface



			EPA Residential	Property 011 Sample ID		Grid 3	Grid 3	Grid 3
			Soil Screening	Depth		2-6in bgs	6-12in bgs	12-24in bgs
Analista	CACNO	l lusita	_	Date	•	_	_	•
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Metals	T	1	1				1	1
Aluminum	7429-90-5	mg/kg	77000		7300	10300 J	8400 J	9990 J
Arsenic	7440-38-2	mg/kg	0.68		1.6 J	1.9	2.1	3.7
Barium	7440-39-3	mg/kg	15000		68.4	62.5 J	96.1 J	180 J
Beryllium	7440-41-7	mg/kg	160		0.54 U	0.57	0.85	1.2
Calcium	7440-70-2	mg/kg	NP		20300 J	2580	2970	2660
Chromium	7440-47-3	mg/kg	NP		11.3 J	14.9	23.4	29.7
Cobalt	7440-48-4	mg/kg	23		3.5	4.2	5.2	8.5
Copper	7440-50-8	mg/kg	3100		6.4 J	6.7	11.2	9.3
Iron	7439-89-6	mg/kg	55000		9820	16100 J	13500 J	17900 J
Lead	7439-92-1	mg/kg	400		21.8	15.4	15.1	6.4
Magnesium	7439-95-4	mg/kg	NP		1720 J	1400	1220	1470
Manganese	7439-96-5	mg/kg	1800		133 J	102	219	323
Mercury	7439-97-6	mg/kg	9.4		0.0081 LJ	0.1 U	0.11 U	0.11 U
Nickel	7440-02-0	mg/kg	1500		7	8.7	13.6	17.2
Potassium	7440-09-7	mg/kg	NP		1210	1100	743	729
Selenium	7782-49-2	mg/kg	390		2.7 UJ	0.31 LJ	2.1 U	2.3 U
Vanadium	7440-62-2	mg/kg	390		16.5	22.5	31.7	41.3
Zinc	7440-66-6	mg/kg	23000		33.9 J	22.5 J	24.1 J	24.8 J



			EPA	Property 011				
			Residential	Sample ID	Grid 3	Grid 3	Grid 3	Grid 3
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Polychlorinated Biphenyls (No	ne Detected)							
Pesticides								
4,4'-DDD	72-54-8	mg/kg	2.3		0.00063 LJ	0.0002 LJ	0.0038 U	0.00024 LJ
4,4'-DDE	72-55-9	mg/kg	2		0.00012 LJ	0.004 U	0.000055 LJ	0.00007 LJ
4,4'-DDT	50-29-3	mg/kg	1.9		0.00018 LJ	0.00021 LJ	0.00011 LJ	0.0038 U
Aldrin	309-00-2	mg/kg	0.039		0.0023 U	0.00029 LJ	0.002 U	0.002 U
alpha-BHC	319-84-6	mg/kg	0.086		0.00012 LJ	0.002 U	0.002 U	0.002 U
alpha-Chlordane	5103-71-9	mg/kg	NP		0.0023 U	0.00041 LJ	0.002 U	0.002 U
beta-BHC	319-85-7	mg/kg	0.3		0.00036 LJ	0.002 U	0.002 U	0.0001 LJ
delta-BHC	319-86-8	mg/kg	NP		0.00014 LJ	0.002 U	0.002 U	0.002 U
Dieldrin	60-57-1	mg/kg	0.034		0.00048 LJ	0.004 U	0.0038 U	0.0038 U
Endosulfan I	959-98-8	mg/kg	NP		0.0023 U	0.002 U	0.002 U	0.002 U
Endosulfan II	33213-65-9	mg/kg	NP		0.00011 LJ	0.000045 LJ	0.00019 LJ	0.000017 LJ
Endosulfan sulfate	1031-07-8	mg/kg	NP		0.00076 LJ	0.000062 LJ	0.000081 LJ	0.0038 U
Endrin	72-20-8	mg/kg	19		0.0044 U	0.00024 LJ	0.0038 U	0.00044 LJ
Endrin aldehyde	7421-93-4	mg/kg	NP		0.0044 U	0.004 U	0.0038 U	0.0038 U
Endrin ketone	53494-70-5	mg/kg	NP		0.0044 U	0.000078 LJ	0.0038 U	0.0038 U
gamma-BHC (Lindane)	58-89-9	mg/kg	0.57		0.00028 LJ	0.002 U	0.002 U	0.002 U
gamma-Chlordane	5103-74-2	mg/kg	NP		0.00026 LJ	0.00058 LJ	0.0002 LJ	0.002 U
Heptachlor	76-44-8	mg/kg	0.13		0.00061 LJ	0.00048 LJ	0.002 U	0.002 U
Heptachlor epoxide	1024-57-3	mg/kg	0.07		0.00063 LJ	0.002 U	0.002 U	0.002 U
Methoxychlor	72-43-5	mg/kg	320		0.00014 LJ	0.02 U	0.02 U	0.02 U



			EPA Residential Soil Screening	Property 011 Sample ID Depth	Grid 3 0-2in bgs	Grid 3 2-6in bgs	Grid 3 6-12in bgs	Grid 3 12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Semi-volatile Organic Compou	ınds							
Bis(2-ethylhexyl)phthalate	117-81-7	mg/kg	39		0.039 LJ	0.019 LJ	0.2 U	0.2 U
Pyrene	129-00-0	mg/kg	1800		NA	NA	NA	NA
Semi-volatile Organic Compou	ınds (SIM)							
2-Methylnaphthalene	91-57-6	mg/kg	240		0.0019 LJ	0.0011 LJ	0.0038 U	0.0039 U
Acenaphthene	83-32-9	mg/kg	3600		0.0044 UJ	0.004 U	0.0038 U	0.0039 U
Acenaphthylene	208-96-8	mg/kg	NP		0.0044 UJ	0.004 U	0.0038 U	0.0039 U
Anthracene	120-12-7	mg/kg	18000		0.0017 LJ	0.002 LJ	0.0038 U	0.0039 U
Benzo(a)anthracene	56-55-3	mg/kg	0.16		0.0063 J	0.0065	0.001 LJ	0.0039 U
Benzo(a)pyrene	50-32-8	mg/kg	0.016		0.0046 J	0.0042	0.0038 U	0.0039 U
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16		0.0078 J	0.0076	0.00096 LJ	0.0039 U
Benzo(g,h,i)perylene	191-24-2	mg/kg	NP		0.0018 LJ	0.0025 LJ	0.0038 U	0.0039 U
Benzo(k)fluoranthene	207-08-9	mg/kg	1.6		0.0021 LJ	0.0018 LJ	0.0038 U	0.0039 U
Chrysene	218-01-9	mg/kg	16		0.0096 J	0.0095	0.0019 LJ	0.0039 U
Dibenzo(a,h)anthracene	53-70-3	mg/kg	0.016		0.0012 LJ	0.0014 LJ	0.0038 U	0.0039 U
Fluoranthene	206-44-0	mg/kg	2400		0.0073 J	0.007	0.00077 LJ	0.0039 U
Fluorene	86-73-7	mg/kg	2400		0.0044 UJ	0.004 U	0.0038 U	0.0039 U
Indeno(1,2,3-cd)pyrene	193-39-5	mg/kg	0.16		0.004 LJ	0.0049	0.00078 LJ	0.0039 U
Naphthalene	91-20-3	mg/kg	3.8		0.0011 LJ	0.00086 LJ	0.0038 U	0.0039 U
Phenanthrene	85-01-8	mg/kg	NP		0.0055 J	0.0038 LJ	0.0013 LJ	0.0039 U
Pyrene	129-00-0	mg/kg	1800		0.0096 J	0.0093	0.0016 LJ	0.0039 U



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	-	Grid 3	Grid 3 2-6in bgs 6/2/2015	Grid 3 6-12in bgs 6/2/2015	Grid 3 12-24in bgs 6/2/2015
Volatile Organic Compounds								
2-Butanone	78-93-3	mg/kg	27000		0.022 U	0.0097 U	0.0093 U	0.0096 U
Acetone	67-64-1	mg/kg	61000		0.0052 LJ	0.0059 LJ	0.0093 U	0.0089 LJ
Ethylbenzene	100-41-4	mg/kg	5.8		0.011 U	0.0049 U	0.0047 U	0.00025 LJ
m,p-Xylene	108-38-3/106-42-3	mg/kg	550		0.011 U	0.00044 LJ	0.00022 LJ	0.00093 LJ
o-Xylene	95-47-6	mg/kg	650		0.00055 LJ	0.0049 U	0.0047 U	0.00031 LJ
Toluene	108-88-3	mg/kg	4900		0.013 J^	0.012	0.0043 LJ	0.02

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL NA - Not Analyzed
Yellow highlighted cells exceed RSL NP - Not Published
U - Not Detected at reported quantitation limit J - Estimated Value
L - Reported concentration is below the CRQL R - Unusable

mg/kg-milligrams per kilogram SIM-selected ion monitoring

BOLD-indicates a detected concentration in-inches

bgs-below ground surface



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	Property 011 Sample ID Depth Date	Grid 4 0-2in bgs 6/1/2015	Grid 4 2-6in bgs 6/2/2015	Grid 4 6-12in bgs 6/2/2015	Grid 4 12-24in bgs 6/2/2015
Metals	CHOING	- Onnes	1000.	2.000	0, 1, 1010	0, 2, 2020	0, 2, 2020	0/ =/ =0=0
Aluminum	7429-90-5	mg/kg	77000		4270	6110 J	12700 J	5260 J
Arsenic	7440-38-2	mg/kg	0.68		1.4	1.5	1.6	1.5
Barium	7440-39-3	mg/kg	15000		51.9	69.3 J	88.4 J	53.7 J
Beryllium	7440-41-7	mg/kg	160		0.43 U	0.5 U	0.62	0.54 U
Calcium	7440-70-2	mg/kg	NP		4150	5130	2700	1540
Chromium	7440-47-3	mg/kg	NP		10	17.7	14.8	11.7
Cobalt	7440-48-4	mg/kg	23		2.7	3.3	6.3	3.4
Copper	7440-50-8	mg/kg	3100		7	8	8.6	5.3
Iron	7439-89-6	mg/kg	55000		7150	9340 J	18100 J	10000 J
Lead	7439-92-1	mg/kg	400		28.6	15	16.2	8
Magnesium	7439-95-4	mg/kg	NP		1030	965	1350	657
Manganese	7439-96-5	mg/kg	1800		158	169	179	105
Mercury	7439-97-6	mg/kg	9.4		0.027 LJ	0.0046 LJ	0.11 U	0.1 U
Nickel	7440-02-0	mg/kg	1500		5.8	9.3	8.9	7.4
Potassium	7440-09-7	mg/kg	NP		704	823	1220	563
Selenium	7782-49-2	mg/kg	390		0.44 LJ	0.38 LJ	0.4 LJ	2.7 U
Vanadium	7440-62-2	mg/kg	390		13.8	27.2	22.6	17.7
Zinc	7440-66-6	mg/kg	23000		21.8	19.6 J	18.8 J	12.9 J



			EPA	Property 011				
			Residential	Sample ID	Grid 4	Grid 4	Grid 4	Grid 4
			Soil Screening	Depth	0-2in bgs	2-6in bgs	6-12in bgs	12-24in bgs
Analyte	CAS.NO	Units	Level	Date	6/1/2015	6/2/2015	6/2/2015	6/2/2015
Polychlorinated Biphenyls (No	ne Detected)							
Pesticides								
4,4'-DDD	72-54-8	mg/kg	2.3		0.00095 LJ	0.0039 U	0.00027 LJ	0.00079 LJ
4,4'-DDE	72-55-9	mg/kg	2		0.00065 LJ	0.000086 LJ	0.00004 LJ	0.00025 LJ
4,4'-DDT	50-29-3	mg/kg	1.9		0.00095 LJ	0.00027 LJ	0.00023 LJ	0.00016 LJ
Aldrin	309-00-2	mg/kg	0.039		0.00079 LJ	0.002 U	0.002 U	0.0019 U
alpha-BHC	319-84-6	mg/kg	0.086		0.0021 U	0.002 U	0.002 U	0.0019 U
alpha-Chlordane	5103-71-9	mg/kg	NP		0.0021 U	0.002 U	0.002 U	0.0019 U
beta-BHC	319-85-7	mg/kg	0.3		0.0021 U	0.000096 LJ	0.002 U	0.00032 LJ
delta-BHC	319-86-8	mg/kg	NP		0.0021 U	0.002 U	0.002 U	0.0019 U
Dieldrin	60-57-1	mg/kg	0.034		0.00054 LJ	0.0039 U	0.0038 U	0.0037 U
Endosulfan I	959-98-8	mg/kg	NP		0.0005 LJ	0.002 U	0.002 U	0.0019 U
Endosulfan II	33213-65-9	mg/kg	NP		0.000027 LJ	0.00066 LJ	0.00026 LJ	0.000042 LJ
Endosulfan sulfate	1031-07-8	mg/kg	NP		0.0041 U	0.00021 LJ	0.0038 U	0.000057 LJ
Endrin	72-20-8	mg/kg	19		0.0041 U	0.00074 LJ	0.00066 LJ	0.0037 U
Endrin aldehyde	7421-93-4	mg/kg	NP		0.00022 LJ	0.0039 U	0.0038 U	0.0037 U
Endrin ketone	53494-70-5	mg/kg	NP		0.0041 U	0.00024 LJ	0.0038 U	0.0037 U
gamma-BHC (Lindane)	58-89-9	mg/kg	0.57		0.00033 LJ	0.002 U	0.002 U	0.0019 U
gamma-Chlordane	5103-74-2	mg/kg	NP		0.0021 U	0.00038 LJ	0.00023 LJ	0.000058 LJ
Heptachlor	76-44-8	mg/kg	0.13		0.00018 LJ	0.002 U	0.002 U	0.0019 U
Heptachlor epoxide	1024-57-3	mg/kg	0.07		0.0021 U	0.002 U	0.002 U	0.0019 U
Methoxychlor	72-43-5	mg/kg	320		0.021 U	0.02 U	0.02 U	0.019 U



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	Property 011 Sample ID Depth Date	Grid 4 0-2in bgs 6/1/2015	Grid 4 2-6in bgs 6/2/2015	Grid 4 6-12in bgs 6/2/2015	Grid 4 12-24in bgs 6/2/2015
Semi-volatile Organic Compou		Offics	Level	Date	0/1/2013	0/2/2013	0/2/2013	0/2/2013
Bis(2-ethylhexyl)phthalate	117-81-7	mg/kg	39		0.024 LJ	0.021 LJ	0.2 U	0.19 U
Pyrene	129-00-0	mg/kg	1800		NA	NA	NA	NA
Semi-volatile Organic Compou	inds (SIM)	<u> </u>						
2-Methylnaphthalene	91-57-6	mg/kg	240		0.0011 LJ	0.0033 LJ	0.0076 U	0.0037 U
Acenaphthene	83-32-9	mg/kg	3600		0.0041 UJ	0.0039 U	0.0076 U	0.0037 U
Acenaphthylene	208-96-8	mg/kg	NP		0.0025 LJ	0.0016 LJ	0.0042 LJ	0.003 LJ
Anthracene	120-12-7	mg/kg	18000		0.0051 J	0.0038 LJ	0.0079	0.0039
Benzo(a)anthracene	56-55-3	mg/kg	0.16		0.012 J	0.0095	0.024	0.013
Benzo(a)pyrene	50-32-8	mg/kg	0.016		0.0092 J	0.0067	0.017	0.0084
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16		0.015 J	0.012	0.033	0.016
Benzo(g,h,i)perylene	191-24-2	mg/kg	NP		0.0093 J	0.0015 LJ	0.0029 LJ	0.0037 U
Benzo(k)fluoranthene	207-08-9	mg/kg	1.6		0.0045 J	0.0033 LJ	0.0089	0.0044
Chrysene	218-01-9	mg/kg	16		0.018 J	0.015	0.037	0.018
Dibenzo(a,h)anthracene	53-70-3	mg/kg	0.016		0.0028 LJ	0.002 LJ	0.0055 LJ	0.0025 LJ
Fluoranthene	206-44-0	mg/kg	2400		0.014 J	0.011	0.027	0.012
Fluorene	86-73-7	mg/kg	2400		0.0041 UJ	0.0039 U	0.0076 U	0.0037 U
Indeno(1,2,3-cd)pyrene	193-39-5	mg/kg	0.16		0.0099 J	0.0066	0.019	0.0084
Naphthalene	91-20-3	mg/kg	3.8		0.0041 UJ	0.002 LJ	0.0076 U	0.0037 U
Phenanthrene	85-01-8	mg/kg	NP		0.0067 J	0.0068	0.0095	0.0055
Pyrene	129-00-0	mg/kg	1800		0.018 J	0.016	0.032	0.015



Analyte	CAS.NO	Units	EPA Residential Soil Screening Level	Property 011 Sample ID Depth Date	Grid 4 0-2in bgs	Grid 4 2-6in bgs 6/2/2015	Grid 4 6-12in bgs 6/2/2015	Grid 4 12-24in bgs 6/2/2015
Volatile Organic Compounds								
2-Butanone	78-93-3	mg/kg	27000		0.012 U	0.011 U	0.01 U	0.012
Acetone	67-64-1	mg/kg	61000		0.01 LJ	0.011 U	0.01	0.0099 LJ
Ethylbenzene	100-41-4	mg/kg	5.8		0.006 U	0.0056 U	0.005 U	0.0051 U
m,p-Xylene	108-38-3/106-42-3	mg/kg	550		0.00049 LJ	0.0056 U	0.00027 LJ	0.00025 LJ
o-Xylene	95-47-6	mg/kg	650		0.006 U	0.0056 U	0.005 U	0.0051 U
Toluene	108-88-3	mg/kg	4900		0.011 J	0.0029 LJ	0.0021 LJ	0.0029 LJ

NA - Not Analyzed

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL

Yellow highlighted cells exceed RSL NP - Not Published

U - Not Detected at reported quantitation limit J - Estimated Value

L - Reported concentration is below the CRQL R - Unusable

mg/kg-milligrams per kilogram SIM-selected ion monitoring

BOLD-indicates a detected concentration in-inches

bgs-below ground surface



-ull List of Chemicals Anlyzed in the Soil

-uii List of Chemicals Anlyzed in the Soi			
			EPA Residential
	GLGNO		Soil Screening
Analyte	CAS.NO	Units	Level
Metals	7420.00.5		77000
Aluminum	7429-90-5	mg/kg	77000
Antimony	7440-36-0	mg/kg	31
Arsenic	7440-38-2	mg/kg	0.68
Barium	7440-39-3	mg/kg	15000
Beryllium	7440-41-7	mg/kg	160
Cadmium	7440-43-9	mg/kg	71
Calcium	7440-70-2	mg/kg	NP
Chromium	7440-47-3	mg/kg	NP
Cobalt	7440-48-4	mg/kg	23
Copper	7440-50-8	mg/kg	3100
Iron .	7439-89-6	mg/kg	55000
Lead	7439-92-1	mg/kg	400
Magnesium	7439-95-4	mg/kg	NP
Manganese	7439-96-5	mg/kg	1800
Mercury	7439-97-6	mg/kg	9.4
Nickel	7440-02-0	mg/kg	1500
Potassium	7440-09-7	mg/kg	NP
Selenium	7782-49-2	mg/kg	390
Silver	7440-22-4	mg/kg	390
Sodium	7440-23-5	mg/kg	NP
Thallium	7440-28-0	mg/kg	0.78
Vanadium	7440-62-2	mg/kg	390
Zinc	7440-66-6	mg/kg	23000
Pesticides			
4,4'-DDD	72-54-8	mg/kg	2.3
4,4'-DDE	72-55-9	mg/kg	2
4,4'-DDT	50-29-3	mg/kg	1.9
Acetophenone	98-86-2	mg/kg	7800
Aldrin	309-00-2	mg/kg	0.039
alpha-BHC	319-84-6	mg/kg	0.086
alpha-Chlordane	5103-71-9	mg/kg	NP
Atrazine	1912-24-9	mg/kg	2.4
Benzaldehyde	100-52-7	mg/kg	7800
beta-BHC	319-85-7	mg/kg	0.3
Caprolactam	105-60-2	mg/kg	31000
Carbazole	86-74-8	mg/kg	NP
delta-BHC	319-86-8	mg/kg	NP
Dieldrin	60-57-1	mg/kg	0.034
Endosulfan I	959-98-8	mg/kg	NP
Endosulfan II	33213-65-9	mg/kg	NP
Endosulfan sulfate	1031-07-8	mg/kg	NP
Endrin	72-20-8	mg/kg	19
Endrin aldehyde	7421-93-4	mg/kg	NP
Endrin ketone	53494-70-5	mg/kg	NP
gamma-BHC (Lindane)	58-89-9	mg/kg	0.57
gamma-Chlordane	5103-74-2	mg/kg	NP
Heptachlor	76-44-8	mg/kg	0.13
Heptachlor epoxide	1024-57-3	mg/kg	0.07
Methoxychlor	72-43-5	mg/kg	320
Toxaphene	8001-35-2	mg/kg	0.49

mg/kg-milligrams per kilogram

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL

			EPA Residential Soil Screening
Analyte	CAS.NO	Units	Level
Polychlorinated Biphenyls	4067444.0	/1	
Aroclor-1016	12674-11-2	mg/kg	4.1
Aroclor-1221 Aroclor-1232	11104-28-2	mg/kg	0.17
Aroclor-1242	11141-16-5 53469-21-9	mg/kg mg/kg	0.17 0.23
Aroclor-1248	12672-29-6	mg/kg	0.23
Aroclor-1254	11097-69-1	mg/kg	0.24
Aroclor-1260	11096-82-5	mg/kg	0.24
Aroclor-1262	37324-23-5	mg/kg	NP
Aroclor-1268	11100-14-4	mg/kg	NP
Volatile Organic Compounds			
1,1,1-Trichloroethane	71-55-6	mg/kg	8100
1,1,2,2-Tetrachloroethane	79-34-5	mg/kg	0.6
1,1,2-Trichloroethane	79-00-5	mg/kg	1.1
1,1-Dichloroethane	75-34-3	mg/kg	3.6
1,1-Dichloroethene	75-35-4	mg/kg	230
1,2,3-Trichlorobenzene	87-61-6	mg/kg	63
1,2-Dibromo-3-chloropropane	96-12-8	mg/kg	0.0053
1,2-Dibromoethane	106-93-4	mg/kg	0.036
1,2-Dichlorobenzene	95-50-1	mg/kg	1800
1,2-Dichloroethane	107-06-2	mg/kg	0.46
1,2-Dichloropropane	78-87-5	mg/kg	1
1,3-Dichlorobenzene	541-73-1	mg/kg	NP
1,4-Dichlorobenzene 2-Butanone	106-46-7 78-93-3	mg/kg	2.6 27000
2-Hexanone	591-78-6	mg/kg	200
4-Methyl-2-Pentanone	108-10-1	mg/kg mg/kg	5300
Acetone	67-64-1	mg/kg	61000
Benzene	71-43-2	mg/kg	1.2
Bromochloromethane	74-97-5	mg/kg	150
Bromodichloromethane	75-27-4	mg/kg	0.29
Bromoform	75-25-2	mg/kg	19
Bromomethane	74-83-9	mg/kg	6.8
Carbon disulfide	75-15-0	mg/kg	770
Carbon tetrachloride	56-23-5	mg/kg	0.65
Chlorobenzene	108-90-7	mg/kg	280
Chloroethane	75-00-3	mg/kg	14000
Chloroform	67-66-3	mg/kg	0.32
Chloromethane	74-87-3	mg/kg	110
cis-1,2-Dichloroethene	156-59-2	mg/kg	160
cis-1,3-Dichloropropene	10061-01-5	mg/kg	NP
Cyclohexane	110-82-7	mg/kg	6500
Dibromochloromethane	124-48-1	mg/kg	0.75
Dichlorodifluoromethane	75-71-8	mg/kg	87
Ethylbenzene 	100-41-4	mg/kg	5.8
Isopropylbenzene	98-82-8	mg/kg	1900
m,p-Xylene Methyl acetate	3-38-3/106-4 79-20-9	mg/kg mg/kg	550 78000
Methyl tert-butyl ether	1634-04-4		47
Methylcyclohexane	108-87-2	mg/kg mg/kg	NP
Methylene chloride	75-09-2	mg/kg	57
o-Xylene	95-47-6	mg/kg	650
Styrene	100-42-5	mg/kg	6000
Tetrachloroethene	127-18-4	mg/kg	24
Toluene	108-88-3	mg/kg	4900
trans-1,2-Dichloroethene	156-60-5	mg/kg	1600
trans-1,3-Dichloropropene	10061-02-6	mg/kg	NP
Trichloroethene	79-01-6	mg/kg	0.94
Trichlorofluoromethane	75-69-4	mg/kg	730
Vinyl chloride	75-01-4	mg/kg	0.059

NP - Not Published

²ull List of Chemicals Anlyzed in the Soil

			EPA Residential
	a.a.va		Soil Screening
Analyte	CAS.NO	Units	Level
Semi-volatile Organic Compounds	76 12 1	ma/ka	40000
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	mg/kg	40000
1,1'-Biphenyl	92-52-4	mg/kg	47
1,2,4,5-Tetrachlorobenzene 1,2,4-Trichlorobenzene	95-94-3	mg/kg	23 24
1,4-Dioxane	120-82-1 123-91-1	mg/kg	5.3
2,2'-Oxybis(1-chloropropane)	108-60-1	mg/kg mg/kg	4.9
2,3,4,6-Tetrachlorophenol	58-90-2	mg/kg	1900
2,4,5-Trichlorophenol	95-95-4	mg/kg	6300
2,4,6-Trichlorophenol	88-06-2	mg/kg	49
2,4-Dichlorophenol	120-83-2	mg/kg	190
2,4-Dimethylphenol	105-67-9	mg/kg	1300
2,4-Dinitrophenol	51-28-5	mg/kg	130
2,4-Dinitrotoluene	121-14-2	mg/kg	1.7
2,6-Dinitrotoluene	606-20-2	mg/kg	0.36
2-Chloronaphthalene	91-58-7	mg/kg	4800
2-Chlorophenol	95-57-8	mg/kg	390
2-Methylnaphthalene	91-57-6	mg/kg	240
2-Methylphenol	95-48-7	mg/kg	3200
2-Nitroaniline	88-74-4	mg/kg	630
2-Nitrophenol	88-75-5	mg/kg	NP
3,3'-Dichlorobenzidine	91-94-1	mg/kg	1.2
3-Nitroaniline	99-09-2	mg/kg	NP
4,6-Dinitro-2-methylphenol	534-52-1	mg/kg	5.1
4-Bromophenyl-phenylether	101-55-3	mg/kg	NP
4-Chloro-3-methylphenol	59-50-7	mg/kg	6300
4-Chloroaniline	106-47-8	mg/kg	2.7
4-Chlorophenyl-phenylether	7005-72-3	mg/kg	NP
4-Methylphenol	106-44-5	mg/kg	6300
4-Nitroaniline	100-01-6	mg/kg	27
4-Nitrophenol	100-02-7	mg/kg	NP
Benzo(a)anthracene	56-55-3	mg/kg	0.16
Benzo(a)pyrene	50-32-8	mg/kg	0.016
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16
Bis(2-chloroethoxy)methane	111-91-1	mg/kg	190
Bis(2-chloroethyl)ether	111-44-4	mg/kg	0.23
Bis(2-ethylhexyl)phthalate	117-81-7	mg/kg	39
Butylbenzylphthalate	85-68-7	mg/kg	290
Chrysene	218-01-9	mg/kg	16
Dibenzofuran	132-64-9	mg/kg	73
Diethylphthalate	84-66-2	mg/kg	51000
Dimethylphthalate	131-11-3	mg/kg	NP
Di-n-butylphthalate	84-74-2	mg/kg	6300
Di-n-octylphthalate	117-84-0	mg/kg	630
Fluoranthene	206-44-0	mg/kg	2400
Hexachlorobenzene	118-74-1	mg/kg	0.21
Hexachlorobutadiene	87-68-3	mg/kg	1.2
Hexachlorocyclopentadiene	77-47-4	mg/kg	1.8
Hexachloroethane	67-72-1 78-50-1	mg/kg	1.8
Isophorone Nitrobonzono	78-59-1	mg/kg	570 E 1
Nitrobenzene	98-95-3	mg/kg	5.1
N-Nitroso-di-n-propylamine	621-64-7	mg/kg	0.078
N-Nitrosodiphenylamine	86-30-6	mg/kg	110
Pentachlorophenol	87-86-5	mg/kg	1
Phenanthrene	85-01-8	mg/kg	NP
Phenol Pyrene	108-95-2 129-00-0	mg/kg mg/kg	19000 1800

			EPA Residential Soil Screening
Analyte	CAS.NO	Units	Level
Semi-volatile Organic Compounds	s (SIM)		
2-Methylnaphthalene	91-57-6	mg/kg	240
Acenaphthene	83-32-9	mg/kg	3600
Acenaphthylene	208-96-8	mg/kg	NP
Anthracene	120-12-7	mg/kg	18000
Benzo(a)anthracene	56-55-3	mg/kg	0.16
Benzo(a)pyrene	50-32-8	mg/kg	0.016
Benzo(b)fluoranthene	205-99-2	mg/kg	0.16
Benzo(g,h,i)perylene	191-24-2	mg/kg	NP
Benzo(k)fluoranthene	207-08-9	mg/kg	1.6
Chrysene	218-01-9	mg/kg	16
Dibenzo(a,h)anthracene	53-70-3	mg/kg	0.016
Fluoranthene	206-44-0	mg/kg	2400
Fluorene	86-73-7	mg/kg	2400
Indeno(1,2,3-cd)pyrene	193-39-5	mg/kg	0.16
Naphthalene	91-20-3	mg/kg	3.8
Pentachlorophenol	87-86-5	mg/kg	1
Phenanthrene	85-01-8	mg/kg	NP
Pyrene	129-00-0	mg/kg	1800

mg/kg-milligrams per kilogram

NP - Not Published

Residential Soil Screening Level- June 2015 EPA Residential Soil RSL

SIM-selected ion monito

Enclosure 4: Agency for Toxic Substances and Disease Registry Fact Sheets

Enclosure 4: Agency for Toxic Substances and Disease Registry fact sheets for Arsenic and Polycyclic Aromatic Hydrocarbons

Enclosure 4: Agency for Toxic Substances and Disease Registry Fact Sheets

Arsenic - ToxFAQs™

CAS # 7440-38-2

This fact sheet answers the most frequently asked health questions (FAQs) about arsenic. For more information, call the CDC Information Center at 1-800-232-4636. This fact sheet is one in a series of summaries about hazardous substances and their health effects. It is important you understand this information because this substance may harm you. The effects of exposure to any hazardous substance depend on the dose, the duration, how you are exposed, personal traits and habits, and whether other chemicals are present.

HIGHLIGHTS: Exposure to higher than average levels of arsenic occur mostly in the workplace, near hazardous waste sites, or in areas with high natural levels. At high levels, inorganic arsenic can cause death. Exposure to lower levels for a long time can cause a discoloration of the skin and the appearance of small corns or warts. Arsenic has been found in at least 1,149 of the 1,684 National Priority List (NPL) sites identified by the Environmental Protection Agency (EPA).

What is arsenic?

Arsenic is a naturally occurring element widely distributed in the earth's crust. In the environment, arsenic is combined with oxygen, chlorine, and sulfur to form inorganic arsenic compounds. Arsenic in animals and plants combines with carbon and hydrogen to form organic arsenic compounds.

Inorganic arsenic compounds are mainly used to preserve wood. Copper chromated arsenate (CCA) is used to make "pressure-treated" lumber. CCA is no longer used in the U.S. for residential uses; it is still used in industrial applications. Organic arsenic compounds are used as pesticides, primarily on cotton fields and orchards.

What happens to arsenic when it enters the environment?

- Arsenic occurs naturally in soil and minerals and may enter the air, water, and land from wind-blown dust and may get into water from runoff and leaching.
- Arsenic cannot be destroyed in the environment.
 It can only change its form.
- Rain and snow remove arsenic dust particles from the air.
- Many common arsenic compounds can dissolve in water. Most of the arsenic in water will ultimately end up in soil or sediment.
- Fish and shellfish can accumulate arsenic; most of this arsenic is in an organic form called arsenobetaine that is much less harmful.

How might I be exposed to arsenic?

- Ingesting small amounts present in your food and water or breathing air containing arsenic.
- Breathing sawdust or burning smoke from wood treated with arsenic.
- Living in areas with unusually high natural levels of arsenic in rock.
- Working in a job that involves arsenic production or use, such as copper or lead smelting, wood treating, or pesticide application.

How can arsenic affect my health?

Breathing high levels of inorganic arsenic can give you a sore throat or irritated lungs.

Ingesting very high levels of arsenic can result in death. Exposure to lower levels can cause nausea and vomiting, decreased production of red and white blood cells, abnormal heart rhythm, damage to blood vessels, and a sensation of "pins and needles" in hands and feet.

Ingesting or breathing low levels of inorganic arsenic for a long time can cause a darkening of the skin and the appearance of small "corns" or "warts" on the palms, soles, and torso.

Skin contact with inorganic arsenic may cause redness and swelling.

Almost nothing is known regarding health effects of organic arsenic compounds in humans. Studies in animals show that some simple organic arsenic



Arsenic

CAS # 7440-38-2

compounds are less toxic than inorganic forms. Ingestion of methyl and dimethyl compounds can cause diarrhea and damage to the kidneys.

How likely is arsenic to cause cancer?

Several studies have shown that ingestion of inorganic arsenic can increase the risk of skin cancer and cancer in the liver, bladder, and lungs. Inhalation of inorganic arsenic can cause increased risk of lung cancer. The Department of Health and Human Services (DHHS) and the EPA have determined that inorganic arsenic is a known human carcinogen. The International Agency for Research on Cancer (IARC) has determined that inorganic arsenic is carcinogenic to humans.

How can arsenic affect children?

There is some evidence that long-term exposure to arsenic in children may result in lower IQ scores. There is also some evidence that exposure to arsenic in the womb and early childhood may increase mortality in young adults.

There is some evidence that inhaled or ingested arsenic can injure pregnant women or their unborn babies, although the studies are not definitive. Studies in animals show that large doses of arsenic that cause illness in pregnant females, can also cause low birth weight, fetal malformations, and even fetal death. Arsenic can cross the placenta and has been found in fetal tissues. Arsenic is found at low levels in breast milk.

How can families reduce the risks of exposure to arsenic?

- If you use arsenic-treated wood in home projects, you should wear dust masks, gloves, and protective clothing to decrease exposure to sawdust.
- If you live in an area with high levels of arsenic in water or soil, you should use cleaner sources of water and limit contact with soil.

 If you work in a job that may expose you to arsenic, be aware that you may carry arsenic home on your clothing, skin, hair, or tools. Be sure to shower and change clothes before going home.

Is there a medical test to determine whether I've been exposed to arsenic?

There are tests available to measure arsenic in your blood, urine, hair, and fingernails. The urine test is the most reliable test for arsenic exposure within the last few days. Tests on hair and fingernails can measure exposure to high levels of arsenic over the past 6-12 months. These tests can determine if you have been exposed to above-average levels of arsenic. They cannot predict whether the arsenic levels in your body will affect your health.

Has the federal government made recommendations to protect human health?

The EPA has set limits on the amount of arsenic that industrial sources can release to the environment and has restricted or cancelled many of the uses of arsenic in pesticides. EPA has set a limit of 0.01 parts per million (ppm) for arsenic in drinking water.

The Occupational Safety and Health Administration (OSHA) has set a permissible exposure limit (PEL) of 10 micrograms of arsenic per cubic meter of workplace air (10 µg/m³) for 8 hour shifts and 40 hour work weeks.

References

Agency for Toxic Substances and Disease Registry (ATSDR). 2007. Toxicological Profile for Arsenic (Update). Atlanta, GA: U.S. Department of Health and Human Services. Public Health Service.

Where can I get more information?

For more information, contact the Agency for Toxic Substances and Disease Registry, Division of Toxicology and Human Health Sciences, 1600 Clifton Road NE, Mailstop F-57, Atlanta, GA 30333.

Phone: 1-800-232-4636

ToxFAQs™ Internet address via WWW is http://www.atsdr.cdc.gov/toxfaqs/index.asp.

ATSDR can tell you where to find occupational and environmental health clinics. Their specialists can recognize, evaluate, and treat illnesses resulting from exposure to hazardous substances. You can also contact your community or state health or environmental quality department if you have any more questions or concerns.

August 2007 Page 2 of 2

Enclosure 4: Agency for Toxic Substances and Disease Registry Fact Sheets

Polycyclic Aromatic Hydrocarbons (PAHs) - ToxFAQs™

This fact sheet answers the most frequently asked health questions (FAQs) about polycyclic aromatic hydrocarbons (PAHs). For more information, call the CDC Information Center at 1-800-232-4636. This fact sheet is one in a series of summaries about hazardous substances and their health effects. This information is important because this substance may harm you. The effects of exposure to any hazardous substance depend on the dose, the duration, how you are exposed, personal traits and habits, and whether other chemicals are present.

SUMMARY: Exposure to polycyclic aromatic hydrocarbons usually occurs by breathing air contaminated by wild fires or coal tar, or by eating foods that have been grilled. PAHs have been found in at least 600 of the 1,430 National Priorities List (NPL) sites identified by the Environmental Protection Agency (EPA).

What are polycyclic aromatic hydrocarbons?

(Pronounced pŏl'ĭ-sī'klĭk ăr'ə-măt'ĭk hī'drə-kar'bənz)

Polycyclic aromatic hydrocarbons (PAHs) are a group of over 100 different chemicals that are formed during the incomplete burning of coal, oil and gas, garbage, or other organic substances like tobacco or charbroiled meat. PAHs are usually found as a mixture containing two or more of these compounds, such as soot.

Some PAHs are manufactured. These pure PAHs usually exist as colorless, white, or pale yellow-green solids. PAHs are found in coal tar, crude oil, creosote, and roofing tar, but a few are used in medicines or to make dyes, plastics, and pesticides.

What happens to PAHs when they enter the environment?

- PAHs enter the air mostly as releases from volcanoes, forest fires, burning coal, and automobile exhaust.
- PAHs can occur in air attached to dust particles.
- Some PAH particles can readily evaporate into the air from soil or surface waters.
- PAHs can break down by reacting with sunlight and other chemicals in the air, over a period of days to weeks.
- PAHs enter water through discharges from industrial and wastewater treatment plants.

- Most PAHs do not dissolve easily in water. They stick to solid particles and settle to the bottoms of lakes or rivers.
- Microorganisms can break down PAHs in soil or water after a period of weeks to months.
- In soils, PAHs are most likely to stick tightly to particles; certain PAHs move through soil to contaminate underground water.
- PAH contents of plants and animals may be much higher than PAH contents of soil or water in which they live.

How might I be exposed to PAHs?

- Breathing air containing PAHs in the workplace of coking, coal-tar, and asphalt production plants; smokehouses; and municipal trash incineration facilities.
- Breathing air containing PAHs from cigarette smoke, wood smoke, vehicle exhausts, asphalt roads, or agricultural burn smoke.
- Coming in contact with air, water, or soil near hazardous waste sites.
- Eating grilled or charred meats; contaminated cereals, flour, bread, vegetables, fruits, meats; and processed or pickled foods.
- Drinking contaminated water or cow's milk.
- Nursing infants of mothers living near hazardous waste sites may be exposed to PAHs through their mother's milk.



Polycyclic Aromatic Hydrocarbons

How can PAHs affect my health?

Mice that were fed high levels of one PAH during pregnancy had difficulty reproducing and so did their offspring. These offspring also had higher rates of birth defects and lower body weights. It is not known whether these effects occur in people.

Animal studies have also shown that PAHs can cause harmful effects on the skin, body fluids, and ability to fight disease after both short- and long-term exposure. But these effects have not been seen in people.

How likely are PAHs to cause cancer?

The Department of Health and Human Services (DHHS) has determined that some PAHs may reasonably be expected to be carcinogens.

Some people who have breathed or touched mixtures of PAHs and other chemicals for long periods of time have developed cancer. Some PAHs have caused cancer in laboratory animals when they breathed air containing them (lung cancer), ingested them in food (stomach cancer), or had them applied to their skin (skin cancer).

Is there a medical test to show whether I've been exposed to PAHs?

In the body, PAHs are changed into chemicals that can attach to substances within the body. There are special tests that can detect PAHs attached to these substances in body tissues or blood. However, these tests cannot tell whether any health effects will occur or find out the extent or source of your exposure to the PAHs. The tests aren't usually available in your doctor's office because special equipment is needed to conduct them.

Has the federal government made recommendations to protect human health?

The Occupational Safety and Health Administration (OSHA) has set a limit of 0.2 milligrams of PAHs per cubic meter of air (0.2 mg/m³). The OSHA Permissible Exposure Limit (PEL) for mineral oil mist that contains PAHs is 5 mg/m³ averaged over an 8-hour exposure period.

The National Institute for Occupational Safety and Health (NIOSH) recommends that the average workplace air levels for coal tar products not exceed 0.1 mg/m³ for a 10-hour workday, within a 40-hour workweek. There are other limits for workplace exposure for things that contain PAHs, such as coal, coal tar, and mineral oil.

Glossary

Carcinogen: A substance that can cause cancer.

Ingest: Take food or drink into your body.

References

Agency for Toxic Substances and Disease Registry (ATSDR). 1995. Toxicological profile for polycyclic aromatic hydrocarbons. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

Where can I get more information?

For more information, contact the Agency for Toxic Substances and Disease Registry, Division of Toxicology and Human Health Sciences, 1600 Clifton Road NE, Mailstop F-57, Atlanta, GA 30333.

Phone: 1-800-232-4636.

ToxFAQs™ Internet address via WWW is http://www.atsdr.cdc.gov/toxfaqs/index.asp.

ATSDR can tell you where to find occupational and environmental health clinics. Their specialists can recognize, evaluate, and treat illnesses resulting from exposure to hazardous substances. You can also contact your community or state health or environmental quality department if you have any more questions or concerns.

September 1996 Page 2 of 2